

* Office hours: Mon & Tue 9-10am in Sc. Ctr. 530 + Mondays 1:30-2:30pm Sc. Ctr. 411.

* Last time: topology on a set X = collection $\mathcal{T} \subset \mathcal{P}(X)$ of open sets st.

(1) $\emptyset, X$ are open (2) $U_i, i \in I$ open $\Rightarrow \bigcup_{i \in I} U_i$ open (3) $U_1, \dots, U_n$ open $\Rightarrow U_1 \cap \dots \cap U_n$ open.

Homeomorphisms: when are two top. spaces "the same"?

Def: X, Y are homeomorphic if there exist continuous maps $f: X \rightarrow Y$ and $g: Y \rightarrow X$ st. $f \circ g = \text{id}_Y, g \circ f = \text{id}_X$.

Equivalently, a homeomorphism $f: X \rightarrow Y$ is a continuous bijection st. f^{-1} continuous
I.e.: a bijection $X \leftrightarrow Y$ under which $\mathcal{T}_X \leftrightarrow \mathcal{T}_Y$.


Remark: a continuous bijection need not be a homeomorphism

Eg: X with 2 topologies, $\mathcal{T}'$ strictly finer than $\mathcal{T}$ ($\mathcal{T}' \supsetneq \mathcal{T}$) $\Rightarrow \text{id}_X: (X, \mathcal{T}') \rightarrow (X, \mathcal{T})$ is a bijection, continuous since $U \in \mathcal{T} \Rightarrow \text{id}^{-1}(U) = U \in \mathcal{T}'$, but not homeo.

• say a metric space (X, d) is bounded if $\text{diam}(X) = \sup \{d(p, q) \mid p, q \in X\} < \infty$

This is not a top. property, eg. $f: (-\frac{\pi}{2}, \frac{\pi}{2}) \rightarrow \mathbb{R}$
 $x \mapsto \tan x$

is a homeomorphism ($\tan$ & $\arctan$ are continuous),
so $\mathbb{R}$ is homeo to $(-\frac{\pi}{2}, \frac{\pi}{2})$ (or any open interval in $\mathbb{R}$)



closed sets: Def: a subset A of a topological space X is closed if $X \setminus A$ is open.

$\triangleq$ subsets can be both closed & open, eg. $\emptyset$ and X , or neither (eg. $[0, 1)$ or $\mathbb{Q}$ in $\mathbb{R}$)

Axioms of open sets imply: $\begin{cases} \bullet \emptyset, X \text{ are closed} \\ \bullet \text{arbitrary intersections of closed sets are closed} \\ \bullet \text{finite unions of closed sets are closed.} \end{cases}$

Def: $A \subset X$ any subset $\Rightarrow$ we define

1) the closure of A : $\bar{A} = \text{smallest closed set containing } A = \bigcap_{\substack{F \supset A \\ F \text{ closed}}} F$
($\bar{A} \supset A$, $\bar{A}$ closed since it's $\cap$ of closed)

2) the interior of A , $\text{int}(A) = \text{largest open set contained in } A$
 $= \bigcup_{U \subset A, U \text{ open}} U$ (open).

3) the boundary of A is $\partial A = \bar{A} - \text{int}(A)$



Ex: $A = [0, 1) \subset \mathbb{R}$, usual top. $\Rightarrow \bar{A} = [0, 1], \text{int}(A) = (0, 1), \partial A = \{0, 1\}$

Rmk: • A is closed iff $\bar{A} = A$, open iff $\text{int}(A) = A$.
 • $\overline{X-A} = X - \text{int}(A)$, $\text{int}(X-A) = X - \bar{A}$. (*)

Def: Say $U \subset X$ is a neighborhood of $p \in X$ if U is open and $p \in U$.



→ Prop: (1) $p \in \text{int}(A)$ iff A contains a neighborhood of p .
 (2) $p \in \bar{A}$ iff every neighborhood of p intersects A nontrivially.

(check this! (1) follows from def^{ns}: $p \in \text{int}(A) \Leftrightarrow \exists U$ open st. $p \in U \subset A$.


(2) follows from (1) + (*): $p \in \bar{A} \Leftrightarrow p \notin \text{int}(X-A) \Leftrightarrow \forall U \ni p$ open, $A \cap U \neq \emptyset$).

Def: Say A is dense if $\bar{A} = X$. (ie. every nonempty open subset of X intersects A nontrivially).

Ex: $\mathbb{Q}$ is dense in $\mathbb{R}$ (for usual topology).

Closed sets & limit points:

Def: $x \in X$ is a limit point of $A \subset X$ if, for every neighborhood $U \ni x$,
 $U \cap (A - \{x\}) \neq \emptyset$



Ex: in $\mathbb{R}_{std}$, 1 is a limit point of $(0,1)$ and of $[0,1]$.

1 is not a limit point of $\{\frac{1}{n}, n \geq 1\} \cup \{0\}$, but 0 is.

Prop: $\bar{A} = A \cup \{\text{limit points of } A\}$.

Pf: $A \subset \bar{A}$ by defⁿ, so enough to consider points not in A .

if $x \notin A$, $\forall U \ni x$ neighborhood, $U \cap (A - \{x\}) = U \cap A$ so $x \in \bar{A}$ iff x limit pt.

x is a limit point if there always $\neq \emptyset$ $\uparrow$ $x \in \bar{A}$ if these always $\neq \emptyset$. $\square$

Conclay: A is closed iff A contains all of its limit points.

• Δ What is the connection between limit points and limits of sequences?

Recall: $\{p_n\}$ sequence in X converges to $p \in X$ if $\forall U$ neighborhood of p ,
 $\exists N$ st. $n \geq N \Rightarrow p_n \in U$.

Fact: $p \in X$, if $\exists \{p_n\}$ sequence in $A \subset X$ st. $p_n \rightarrow p$ then $p \in \bar{A}$
 if $\exists \{p_n\}$ seq. in A , $p_n \neq p$ for ∞ many n , $p_n \rightarrow p$, then p is a limit pt of A .

Pf: any neighborhood $U \ni p$ contains p_n for all large n , hence contains points of A .
 (distinct from p in 2nd case) $\square$

The converse is true in metric spaces: if $p \in \bar{A}$ (resp. a limit point of A) then $\textcircled{3}$
 $\forall n > 0 \exists p_n \in B_{1/n}(p) \cap A$ (resp. with $p_n \neq p$), so $\exists$ sequence in A st. $p_n \rightarrow p$.

This holds more generally in spaces whose points have countable bases of neighborhoods
 $U_1 \supset U_2 \supset \dots$ (ie. $\forall p \exists$ hdd $U_1, U_2, \dots$ st. $\forall$ hdd $U \ni p, \exists n$ st. $p \in U_n \subset U$), but not in arbitrary topological spaces!

Hausdorff spaces: In a metric space, a sequence converges to at most one limit.

This is not true in an arbitrary topological space!

Ex: $X = \mathbb{R}$ with finite complement topology: open subsets = $\emptyset$ and $\mathbb{R} - \{\text{finite sets}\}$

let $a_1, a_2, \dots$ be a sequence in X with all a_i distinct.

Then $\forall x \in X$, every neighborhood $U \ni x$ contains all but finitely many of the a_i ,
hence $\exists N$ st. $a_n \in U \forall n \geq N$. Thus the sequence converges to every point of X !


To avoid such pathological behavior:

Def: A top-space is Hausdorff (or T_2) if $\forall x_1 \neq x_2 \in X, \exists$ neighborhoods $U_1 \ni x_1, U_2 \ni x_2$
st. $U_1 \cap U_2 = \emptyset$.

Ex: 1) any metric space is Hausdorff:

given $x_1 \neq x_2$, choose $0 < \varepsilon < \frac{1}{2} d(x_1, x_2)$

then $U_i = B_\varepsilon(x_i)$ disjoint neighborhoods of x_i .



2) the finite complement topology on $\mathbb{R}$ is not Hausdorff, since any two non-empty open sets intersect (in infinitely many points).

3) the discrete topology is always Hausdorff ($U_i = \{x_i\}$ disjoint neighborhoods of x_i)

4) One can show: X Hausdorff, $Y \subset X \Rightarrow$ the subspace top. is Hausdorff.

X, Y Hausdorff $\Rightarrow X \times Y$ Hausdorff. (Homework!)

Thm: if X is Hausdorff then every sequence in X converges to at most one limit.

Proof: assume $x_1, x_2, \dots$ converges to $x \in X$, and let $y \neq x$. Choose $U_x \ni x, U_y \ni y$
disjoint neighborhoods. Since $x_n \rightarrow x, \exists N$ st. $\forall n \geq N x_n \in U_x$.

Hence $x_n \notin U_y$ for $n \geq N$, so the sequence doesn't converge to y . $\square$

Rmk: There's in fact a whole hierarchy of "separation axioms": eg. a weaker one is:

A top-space is T_1 if $\forall x \neq y \in X, \exists U_y \ni y$ neighborhood st. $x \notin U_y$.

equivalently: X is $T_1 \iff \{x\}$ is closed in $X \forall x \in X$. (exercise!)


Munkres $(T_2) \Rightarrow T_1$, but eg. $(\mathbb{R}, \text{finite complement top.})$ is T_1 but not Hausdorff. (4)

- Hausdorff spaces are fairly nice to work with, and we will generally be working with this assumption. There are more subtle reasons why not every Hausdorff topology comes from a metric, but one can give pretty good criteria for a topology to be metrizable involving further separation conditions ("normal" or T_4) (+ a countability condition). We'll see the Urysohn metrization theorem.

Manifolds & CW complexes:

Metric spaces are nice, but they can still be pretty nasty. (We'll see conditions such as local connectedness, local compactness etc. come up). Algebraic topologists like to focus on even nicer spaces. For example:

Def: An n -dimensional topological manifold is a top. space X st. every point $p \in X$ has a neighborhood homeomorphic to $\mathbb{R}^n$ (or equivalently, an open ball in $\mathbb{R}^n$).

Example:  $S^1 \subset \mathbb{R}^2$ is a 1-d. top. manifold;   $\subset \mathbb{R}^3$ 2d top. manifolds

Example:  isn't a top. manifold (vertex looks wrong) - but it is part of a more general class of spaces called CW complexes, built by attaching "cells" (closed balls of dim. 0, 1, ...) onto each other inductively.

We'll see more on this later when we get to alg. top. In decreasing order of generality:
 $\{\text{top spaces}\} \supset \{\text{Hausdorff}\} \supset \{\text{metrizable}\} \supset \{\text{CW-complex}\} \supset \{\text{manifold}\}.$

Topologies on infinite products: given topological spaces X_i , $i \in I$ index set:

What is the natural topology on $X = \prod_{i \in I} X_i = \{(p_i)_{i \in I} \mid p_i \in X_i \forall i \in I\}$?

First idea: Def: the box topology on $\prod_{i \in I} X_i$ has basis $\{\prod_{i \in I} U_i \mid U_i \subset X_i \text{ open } \forall i\}$
(ie. open set = unions of such "boxes")

(this is a basis: $\text{box} \cap \text{box} = \text{box}$, since $(\prod U_i) \cap (\prod V_i) = \prod (U_i \cap V_i)$)

This is actually too fine for most purposes - next time we'll do better.

Example: consider the diagonal map $\Delta: \mathbb{R} \rightarrow \mathbb{R}^\omega = \mathbb{R}^\mathbb{N} (= \mathbb{R}_0 \times \mathbb{R}_1 \times \mathbb{R}_2 \times \dots)$
 $\Delta(x) = (x, x, x, \dots)$

giving $\mathbb{R}^\mathbb{N}$ the box topology, Δ is not continuous!! (unlike case of finite products)

Indeed, let $U = (-1, 1) \times (-\frac{1}{2}, \frac{1}{2}) \times (-\frac{1}{3}, \frac{1}{3}) \times \dots$ open in box topology.

$\Delta^{-1}(U) = \bigcap_{n \geq 1} (-\frac{1}{n}, \frac{1}{n}) = \{0\}$ not open in $\mathbb{R}$.