STD testing







DISCLAIMER: These recommendations are based on those from the Centers for Disease Control and Prevention and do not substitute an appointment with a physician. Based on your sexual orientation, gender identity and lifestyle, you may benefit from more frequent testing. Consult your physician for an STI testing plan that's right for you.

	Who?	What?	When?
	Anyone who has unprotected sex or shares injection drug equipment	HIV	After having unprotected sex or sharing injection drug equipment
	Everyone ages 13 to 64	HIV	At least once in their lifetime
	Sexually active women ages 25 and younger	Gonorrhea Chlamydia	Once a year
IHS) nefit	Sexually active women ages 25 and older with new or multiple sex partners, or a sex partner who has an STI	Gonorrhea Chlamydia HIV Trichomoniasis Hepatitis B Hepatitis C	Once a year If positive, retest 3 months after treatment
	All pregnant women	Syphilis HIV Hepatitis B If younger than 25: Gonorrhea Chlamydia	First prenatal visit At 3 months In third trimester
	Sexually active men who have sex with men Those who have multiple or anonymous partners	Syphilis Chlamydia Gonorrhea HIV If born before 1945: Hepatitis B Hepatitis C	At least once a year Every 3 to 6 months, or more frequently as directed by a physician



	or shares injection drug equipment	
	Everyone ages 13 to 64	HIV
Where can students get tested?	Sexually active women ages 25 and younger	Gonorrhea Chlamydia
STI testing at Harvard If you are on your parents' health insurance or the Student Health Insurance Plan (SHIP), Harvard University Health Services (HUHS) provides STI testing without any out-of-pocket cost to you. If you are on a different health insurance plan, please refer to your benefit guide to confirm coverage for STI testing. To schedule an STI test at HUHS please call: 617-495-5711	Sexually active women ages 25 and older with new or multiple sex partners, or a sex partner who has an STI	Gonorrhea Chlamydia HIV Trichomoniasis Hepatitis B
STI testing within the Boston area	All pregnant women	Hepatitis C Syphilis
If you do not feel comfortable going to HUHS for STI testing, <u>the CDC website</u> has a list of locations for HIV and STI testing in the Cambridge and Boston area and beyond, just enter your zip code.	An pregnane women	HIV Hepatitis B
How often should you get tested? Frequency of testing will vary by each individual person. <u>Check out the CDC's recommendations on how often to get tested.</u>		lf younger than 2 Gonorrhea Chlamydia
	Sexually active men who have sex with men	Syphilis Chlamydia
	Those who have multiple	Gonorrhea HIV