

Stone-Weierstrass theorem:

Thm: Polynomials are dense in $C^0([a,b])$, ie. $\forall f \in C^0([a,b]) \exists P_n$ polynomials (Weierstrass) st. $P_n \rightarrow f$ uniformly on $[a,b]$.

Pf. uses convolution and its use to approximate/smooth functions.

Def: convolution: $(f * g)(x) = \int_{s+t=x} f(s) g(t) dt = \int_{-\infty}^{\infty} f(x-t) g(t) dt = \int_{-\infty}^{\infty} f(s) g(x-s) ds$.

well-def'd if e.g. f and g are (piecewise) C^0 + one of them is compactly supported (ie. 0 outside some $[-M,M]$). (to avoid improper integrals).

Principle: " $f * g$ inherits the best properties of f and g ".

This is because $\|f * g\|_{\infty} \leq \|f\|_{L_1} \|g\|_{\infty}$. where $g_h(t) := g(t+h)$

here (*) $(f * g)(x+h) - (f * g)(x) = \int f(x-t)(g(t+h) - g(t)) dt \leq \|f\|_{L_1} \|g_h - g\|_{\infty}$

\rightarrow If g is C^0 then (over relevant intervals, per compact support)
uniform continuity (over a compact interval) gives $\lim_{h \rightarrow 0} \|g_h - g\|_{\infty} = 0$.
($|g(t+h) - g(t)| < \varepsilon \ \forall t$ when $|h| < \delta$). \Rightarrow continuity of $f * g$.

\rightarrow If g is C^1 (continuously differentiable) then
dividing (*) by h and using mean value thm + unif. continuity of g' on a
compact interval $\Rightarrow f * g$ is continuously differentiable and $(f * g)' = f * g'$.

Hence if g is C^∞ then $f * g$ is C^∞ !! (even if f isn't even continuous)

\rightarrow and... if g is a polynomial of degree d then so is $f * g$!

e.g. because $g^{(d+1)} = 0$ so $(f * g)^{(d+1)} = f * g^{(d+1)} = 0$, or more directly: $g(x) = \sum_{k=0}^d a_k x^k$
 $\Rightarrow (f * g)(x) = \sum_{k=0}^d a_k \int f(t) (x-t)^k dt = \sum_{k=0}^d \sum_{l=0}^k (-1)^l \binom{k}{l} a_k x^{k-l} \underbrace{\int f(t) t^l dt}_{\text{constant}}$
 manifestly a polynomial in x .

Approximate identities:

Def: A sequence of functions $k_n \geq 0$ approximates identity if $\begin{cases} \cdot \int k_n dx = 1 \\ \cdot \forall \delta > 0, \int_{|x| \geq \delta} k_n dx \rightarrow 0 \end{cases}$

Thm: $\left. \begin{array}{l} f \text{ compactly supported \& continuous} \\ k_n \text{ approximate identity} \end{array} \right\} \Rightarrow f * k_n \rightarrow f \text{ uniformly.}$

Pf: $(f * k_n)(x) - f(x) = \int (f(x-t) - f(x)) k_n(t) dt = \int_{|t| \leq \delta} + \int_{|t| \geq \delta}$. Estimate each term as follows.

Given $\varepsilon > 0$, uniform continuity of f on its support $\Rightarrow \exists \delta$ (indep. of x) st. (2)

$$|t| < \delta \Rightarrow |f(x-t) - f(x)| < \frac{\varepsilon}{2}$$

Then $\left| \int_{-\delta}^{\delta} (f(x-t) - f(x)) K_n(t) dt \right| \leq \frac{\varepsilon}{2} \int_{-\delta}^{\delta} K_n(t) dt \leq \frac{\varepsilon}{2}$. (indep. of x).

while $\left| \int_{|t| \geq \delta} (f(x-t) - f(x)) K_n(t) dt \right| \leq 2 \|f\|_{\infty} \int_{|t| \geq \delta} K_n(t) dt \rightarrow 0$ as $n \rightarrow \infty$


becomes $< \frac{\varepsilon}{2}$ for n sufftly large.

$\rightarrow \exists N$ st. $\forall x$, $|f \ast K_n(x) - f(x)| < \varepsilon \quad \forall n \geq N$. n \geq N (indep. of x !). □


Ex: $K_n(x) = c_n (1-x^2)^n$ for $|x| \leq 1$, 0 elsewhere

where $c_n > 0$ is chosen so that $\int_{-1}^1 K_n dx = 1$.

Claim: K_n approximate identity.



Pf: • for $|x| \leq \frac{1}{\sqrt{2n}}$, $(1-x^2)^n \geq 1-nx^2 \geq \frac{1}{2}$, so $\int_{-1}^1 (1-x^2)^n dx \geq \int_{-\frac{1}{\sqrt{2n}}}^{\frac{1}{\sqrt{2n}}} \frac{1}{2} dx = \frac{1}{\sqrt{2n}}$



$$\Rightarrow c_n \leq \sqrt{2n}$$

• for $|x| \geq \delta$, $(1-x^2)^n \leq (1-\delta^2)^n$ so $\int_{|x| \geq \delta} K_n dx \leq 2c_n (1-\delta^2)^n \leq 2\sqrt{2n} (1-\delta^2)^n \xrightarrow{n \rightarrow \infty} 0$

\rightarrow Thm. (Weierstrass):

$\forall f \in C^0([a,b]) \quad \exists p_n$ polynomials st. $p_n \rightarrow f$ uniformly.

Pf: • by linear change of variables, we can assume $[a,b] = [0,1]$.
 • subtracting a degree 1 polynomial from f we can assume $f(0) = f(1) = 0$. Then extend f to \mathbb{R} by $f(x) = 0$ if $x \notin [0,1]$.
 • now let $K_n(x) =$ as above, and $p_n = f \ast K_n$.

Then K_n approx. identity, f C^0 compactly supported $\Rightarrow p_n \rightarrow f$ uniformly.

• p_n is a polynomial of degree $2n$ on $[0,1]$ because, given that $f=0$ outside $[0,1]$, the formula $(f \ast K_n)(x) = \int f(x-t) K_n(t) dt$ for $x \in [0,1]$ doesn't involve the values of K_n outside $[-1,1]$, and $K_n|_{[-1,1]}$ is polynomial. □

• Stone's theorem generalizes this to other families of functions.

Def: $A \subset C^0(K)$ is an algebra if $f, g \in A \Rightarrow f+g \in A$, $cf \in A$, $fg \in A$.
 A separates points if $\forall a \neq b \in K$, $\exists f, g \in A$ st. $f(a) = 1, f(b) = 0$, $g(a) = 0, g(b) = 1$

(0,1 are arbitrary - this is equiv't to $A \rightarrow \mathbb{R}^2$, $f \mapsto (f(a), f(b))$ is surjective $\forall a \neq b$). □

- * For complex-valued functions, further assume it is conjugation-invariant, ie. $f \in A \Rightarrow \bar{f} \in A$ (equivalently: $\operatorname{Re} f \in A, \operatorname{Im} f \in A$). (3)

Thm (Stone): \parallel K compact metric space, $A \subset C^0(K)$ algebra which separates points (+ conjugation invariant in \mathbb{C} case), then A is dense in $(C^0(K), \| \cdot \|_\infty)$ (Weierstrass = special case $K = [a, b]$, A = polynomials).

- Pf:
- \overline{A} (uniform closure of A) is an algebra ($f_n \rightarrow f, g_n \rightarrow g \Rightarrow f+g = \lim(f_n+g_n), fg = \lim(f_n g_n)$) so enough to show assumptions + A closed $\Rightarrow \overline{A} = C^0(K)$
 - given $f \in A$, A algebra & closed $\Rightarrow P(f) \in A$ $\forall P$ polynomial s.t. $P(0) = 0$. By Weierstrass, $|x|$ is a uniform limit of polynomials on $[-M, M]$, so $|f| \in \overline{A} = A$. Hence: $f, g \in A \Rightarrow \max(f, g) = \frac{f+g+|f-g|}{2} \in A$, same for $\min(f, g)$.
 - Now: given $f \in C^0(K)$, $\varepsilon > 0$, want to show $\exists h \in A$ s.t. $\sup |h-f| \leq \varepsilon$. ($\Rightarrow f \in \overline{A} = A$).
given $x \in K \quad \forall y \neq x \quad \exists g_y \in A$ s.t. $\begin{cases} g_y(x) = f(x) & (A \text{ separates points}) \\ g_y(y) = f(y). \end{cases}$
 $\exists U_y \ni y$ open s.t. $g_y > f - \varepsilon$ on U_y ; and K compact $\Rightarrow \exists y_1, \dots, y_n$ s.t. $U_{y_1} \cup \dots \cup U_{y_n} = K$. Then $h_x := \max(g_{y_1}, \dots, g_{y_n}) \in A$ satisfies $\begin{cases} h_x > f - \varepsilon \text{ everywhere} \\ h_x(x) = f(x). \end{cases}$
By the same argument, $\exists x_1, \dots, x_n$ s.t. $k = \min(h_{x_1}, \dots, h_{x_n})$ satisfies $|k-f| < \varepsilon$ everywhere.
 $(\exists V_x \ni x \text{ open s.t. } h_x < f + \varepsilon \text{ on } V_x; K \text{ compact} \Rightarrow \exists x_1, \dots, x_n \text{ s.t. } V_{x_1} \cup \dots \cup V_{x_n} = K)$ \square

Fourier series: we consider continuous 2π -periodic functions $f: \mathbb{R} \rightarrow \mathbb{C}$ with complex values, or equivalently functions on $S^1 \cong \mathbb{R}/2\pi\mathbb{Z}$, with L^2 inner product $\langle f, g \rangle = \frac{1}{2\pi} \int_0^{2\pi} \bar{f}(x)g(x) dx$

The complex exponentials $e_n(x) = e^{inx}$, $n \in \mathbb{Z}$ satisfy $\langle e_i, e_j \rangle = \delta_{ij}$ - orthonormality.

Def. \parallel The Fourier coefficients of f are $c_n(f) = \langle e_n, f \rangle = \frac{1}{2\pi} \int_0^{2\pi} e^{-inx} f(x) dx$.

\rightarrow the Fourier series of f is $\sum_{n \in \mathbb{Z}} c_n e_n = \sum_{-\infty}^{\infty} c_n(f) e^{inx}$

Q: (Fourier, Dirichlet, Fejér, ...) does the Fourier series accurately represent f ?
(e.g. does it converge? to f ?).

Def: Trigonometric polynomials = the vector space of finite linear combinations of e_n .

- * Clearly this is an algebra, complex conj. invariant, and separates points of S^1 , which is compact: hence by Stone-Weierstrass, trig. polynomials are dense in $(C^0(S^1), \| \cdot \|_\infty)$
... hence also in L^2 -norm ($\|f\|_{L^2} = \left(\frac{1}{2\pi} \int |f|^2 dx\right)^{1/2} \leq \sup |f|$).

- * The n^{th} Fourier sum $f_n = s_n(f) = \sum_{-n}^n c_k e^{ikx} = \sum_{-n}^n \langle e_k, f \rangle e_k$ (4)
- is the orthogonal projection of f onto $V_n = \text{span}(e_{-n}, \dots, e_n)$ for $\langle \cdot, \cdot \rangle$.
- Indeed: $\langle e_j, f_n \rangle = \sum_{k=-n}^n c_k \langle e_j, e_k \rangle = c_j = \langle e_j, f \rangle$, so $\langle e_j, f - f_n \rangle = 0 \quad \forall -n \leq j \leq n$.
- Thus: $\forall g \in V_n, \|f - f_n\|_{L^2} \leq \|f - g\|_{L^2}$ - the point of V_n closest to f for $\|\cdot\|_{L^2}$
- (This follows from $(f - f_n) \perp V_n$: $(f - g) = (\underbrace{f - f_n}_{\perp V_n}) + (\underbrace{f_n - g}_{\in V_n}) \Rightarrow \|f - g\|^2 = \|f - f_n\|^2 + \|f_n - g\|^2 \geq \|f - f_n\|^2$)

Theorem: (Parseval) Let $f \in C^0(S^1)$, $c_n = \langle e_n, f \rangle$ Fourier coeffs, $f_n = \sum_{-n}^n c_k e_k$ partial sums.

(1) $f_n \rightarrow f$ in L^2 , i.e. $\|f_n - f\|_{L^2}^2 = \frac{1}{2\pi} \int |f(x) - f_n(x)|^2 dx \rightarrow 0$ as $n \rightarrow \infty$.

(2) $\sum_{n \in \mathbb{Z}} |c_n|^2 = \|f\|_{L^2}^2 = \frac{1}{2\pi} \int_0^{2\pi} |f(x)|^2 dx$ (in particular $\sum |c_n|^2$ converges $\Rightarrow c_n \rightarrow 0$ as $|n| \rightarrow \infty$)

Pf: (1) Since trig. polynomials = $\bigcup_n V_n$ are dense in $(C^0(S^1), \|\cdot\|_{L^2})$,

$\forall \varepsilon > 0 \exists N$ st. $\exists g \in V_N$ with $\|f - g\|_{L^2} < \varepsilon$.

Now for $n \geq N$, $g \in V_N \subset V_n$ and $f_n =$ closest point to f , so

$\|f - f_n\|_{L^2} \leq \|f - g\|_{L^2} < \varepsilon$. This shows $f_n \rightarrow f$ in L^2 .

(2) since $f_n \in V_n$ and $f - f_n \in V_n^\perp$, $\|f\|_{L^2}^2 = \|f_n\|_{L^2}^2 + \|f - f_n\|_{L^2}^2$
 where $\|f_n\|_{L^2}^2 = \left\| \sum_{-n}^n c_k e_k \right\|^2 = \sum_{-n}^n |c_k|^2$ by orthonormality, and
 $\|f - f_n\|_{L^2}^2 \rightarrow 0$ by the first part. \square

Corollary: if $f, g \in C^0(S^1)$ have same Fourier series then $\frac{1}{2\pi} \int |f - g|^2 dx = \sum |c_n(f) - c_n(g)|^2 = 0$, hence $f = g$.

* The fact that $f_n \rightarrow f$ in L^2 is the best approximation (in L^2 norm) of f by trig. polynomials, and that trig. polynomials are dense in $\|\cdot\|_\infty$ (so \exists trig. polynomials $\rightarrow f$ uniformly) makes one hope that $f_n \rightarrow f$ uniformly or at least pointwise... alas not!

Fact: $\exists f \in C^0(S^1)$ st. the Fourier series of f does not converge ($s_n(f)(0)$ unbounded!) (but the example is hard to construct).

Thm (Dirichlet) if f is C^1 then $f_n = s_n(f) \rightarrow f$ uniformly.

The proof uses convolution - redefine, for periodic functions, $(f * g)(x) = \frac{1}{2\pi} \int_0^{2\pi} f(t) g(x-t) dt$.

& note $c_n e_n(x) = \frac{1}{2\pi} \left(\int f(t) e^{-int} dt \right) e^{inx} = (f * e_n)(x)$.

So: $s_n(f) = \sum_{-n}^n c_k e_k = f * \left(\sum_{-n}^n e_k \right) = f * D_n$ where

$$D_n(x) = \sum_{-n}^n e^{ikx} = \frac{e^{i(n+\frac{1}{2})x} - e^{-i(n+\frac{1}{2})x}}{e^{ix/2} - e^{-ix/2}} = \frac{\sin(n+\frac{1}{2})x}{\sin(\frac{x}{2})} \quad \text{Dirichlet kernel} \quad (5)$$

Dirichlet's proof studies this convolution for $f \in C'$ to prove uniform convergence.

The fact that convergence can sometimes fail makes it remarkable that $\forall f \in C^0$, f can be recovered from the partial sums $S_n(f) = f_n = \sum_{-n}^n c_k e^{ikx} \dots$

Thm (Fejér): || If $f \in C^0(S')$ then $\frac{S_0(f) + \dots + S_{n-1}(f)}{n}$ converges uniformly to f .

The reason is that this process amounts to convolution with the Fejér kernel $F_n = \frac{D_0 + \dots + D_{n-1}}{n}$, which actually approximates identity (in the sense seen above) unlike D_n .